



Are you ready for a different approach to your Parkinson's disease?

VYALEV is a 24-hour, continuous delivery of levodopa-based therapy—morning, day, and night.

USE

VYALEV is a prescription medicine used for treatment of advanced Parkinson's disease in adults. VYALEV contains two medicines, foscarnidopa and foslevodopa.

SELECT SAFETY INFORMATION

Do not use VYALEV if you take or have taken a nonselective monoamine oxidase (MAO) inhibitor within the last 14 days. Ask your healthcare provider if you are not sure if you take an MAO inhibitor.

Please see Full Important Safety Information on pages 12-13 and Full Prescribing Information, including Medication Guide.

Are your levodopa pills **not working as well** as they used to?



As Parkinson's disease progresses, it can slow digestion, causing levodopa pills to sit in your stomach longer without being broken down.



Then, once your levodopa medication has been broken down and reaches your small intestine, it has to compete with certain nutrients from the food you eat to get into your bloodstream.

All of this can reduce the amount of levodopa that reaches your brain when you take pills.

Knowing when you're "On" and "Off"



"On" time is when your medication is working well. This means you're within what's called the "sweet spot."




"Off" time is when your medication is not working well or is no longer helping with mobility, slowness, stiffness, and tremor. Every patient's experience with "Off" time is different, and you may experience symptoms unique to you.

As Parkinson's advances, it's harder to stay in the "sweet spot"


Over time, the treatment window (or "sweet spot") gets smaller, and your levodopa levels can more often become too high (resulting in dyskinesia*) or too low (resulting in "Off" time).

When treating advanced Parkinson's with levodopa pills, levels may become too low, or too high.

The goal of a continuously delivered levodopa therapy is to keep levodopa levels within the "sweet spot." This can help smooth the "peaks and valleys" that may be associated with taking pills.



***Dyskinesia:** Bothersome extra movements you don't have control over.



Clinical trial patients taking VYALEV achieved almost 2 more hours of “On” time each day than patients treated with pills alone.

Now you can _____ **more***

More “On” time is possible with 24-hour, continuous levodopa-based therapy

SELECT SAFETY INFORMATION

VYALEV can cause serious side effects, such as falling asleep without warning during normal daily activities; seeing, hearing, or feeling things that are not real; unusual urges; infusion site reactions and infections; new or worsened uncontrolled sudden movements; heart attack or other heart problems (increased blood pressure, fast or irregular heartbeat, or chest pain); and **worsening of the increased pressure in your eyes** (glaucoma).

Please see Full Important Safety Information on pages 12-13 and Full Prescribing Information, including Medication Guide.

VYALEV delivers 24 hours of continuous levodopa-based therapy—morning, day, and night



VYALEV™
foscarbidopa/foslevodopa
Injection for subcutaneous use
12 mg/240 mg per mL




Continuous delivery

VYALEV delivers a levodopa-based therapy under the skin, directly into the body




No dietary limitations

What you eat won't affect your levodopa levels



The VYALEV pump attaches to your skin with a single cannula*



A qualified healthcare team will train you on the use of VYALEV

*A cannula is a soft, flexible tube, less than a millimeter in diameter, inserted into the skin.

SELECT SAFETY INFORMATION

Seeing, hearing, or feeling things that are not real (hallucination), infusion site reactions and infections, and new or worsened uncontrolled sudden movements (dyskinesia) are common and sometimes serious side effects.

Please see Full Important Safety Information on pages 12-13 and [Full Prescribing Information](#), including [Medication Guide](#).

Benefits of VYALEV



VYALEV™
foscarbidopa/foslevodopa
Injection for subcutaneous use
12 mg/240 mg per mL



Superior “On” time vs pills

Almost 2 more hours of “On” time per day without troublesome dyskinesia than treatment with pills alone*



24-hour, continuous delivery of levodopa-based therapy—morning, day, and night



Does NOT require surgery



Replaces all levodopa pills† to free you from your oral levodopa schedule

*VYALEV was studied against oral immediate-release carbidopa/levodopa pills. At week 12, patients taking VYALEV achieved 2.72 hours more “On” time vs 0.97 hours more “On” time for those taking levodopa pills.

†VYALEV replaces levodopa-containing medications and COMT inhibitors. You can take other Parkinson’s disease medications with VYALEV. Keep a supply of oral immediate-release carbidopa/levodopa tablets with you in case you are unable to use VYALEV infusion.

COMT=catechol-O-methyltransferase.

SELECT SAFETY INFORMATION

Suddenly stopping or changing the VYALEV dose may cause withdrawal symptoms, such as fever, confusion, or severe muscle stiffness. Please see Full Important Safety Information on pages 12-13 and [Full Prescribing Information](#), including [Medication Guide](#).

Could VYALEV be right for you?

- ☐ Are you experiencing more than **2½ hours** of “Off” time each day?
- ☐ Are you taking more levodopa pills or **taking them more often**?

If you answered yes to either of these questions, you’re not alone. Talk to your doctor about whether VYALEV may help.

Only you and your doctor can decide if a treatment is right for you. Your doctor is your best resource for medical information.

VYALEVTM
foscarbidopa/foslevodopa
Injection for subcutaneous use
12 mg/240 mg per mL

“VYALEV works for me because it allows me to have **medicine always available when I need it**. I don’t have to worry about whether I’m getting the right dosage, whether I’m ‘On’ or ‘Off.’”

—Janice, a VYALEV patient*

*Janice is an actual VYALEV patient and was on prescribed therapy when she provided a testimonial. Changes in therapy status may have occurred since that time.

SELECT SAFETY INFORMATION

Infusion site infections and reactions could include skin changes at the infusion site, such as redness, warmth, swelling, pain, or discoloration. Remove your cannula and call your healthcare provider if you see any of these signs.

Please see Full Important Safety Information on pages 12-13 and Full Prescribing Information, including Medication Guide.

Designed to fit into your life

VYALEV is a 24-hour, continuous delivery of levodopa-based therapy.



VYALEV™
foscarbidopa/foslevodopa
Injection for subcutaneous use
12 mg/240 mg per mL



**Small and
less than 1 lb (0.63 lb)**

**Large buttons
and clear display**

Can be temporarily disconnected for up
to 1 hour for water-based activities, such as
swimming or showering



**VYALEV and its accessories
can be worn under clothing**

Do NOT use the VYALEV delivery system while engaging in (or performing) any activity that could result in liquids getting onto or into the pump. Some of these activities include taking a bath or shower, or swimming.

Keep a supply of oral immediate-release carbidopa/levodopa tablets with you in case you are unable to use VYALEV infusion. Please refer to the Pump Instructions for Use that will come from your Specialty Pharmacy.

SELECT SAFETY INFORMATION

Do not use VYALEV if you take or have taken a nonselective monoamine oxidase (MAO) inhibitor within the last 14 days. Ask your healthcare provider if you are not sure if you take an MAO inhibitor.

Please see Full Important Safety Information on pages 12-13 and Full Prescribing Information, including Medication Guide.


Follow the 5 steps to administer VYALEV

The **5 Ps (Plan, Prepare, Prime, Place, and Pump)** teach you how to take VYALEV. These steps will help prepare you to administer the product.

A qualified healthcare provider will train you and make sure you are comfortable with the steps of the 5 Ps before you start VYALEV.



VYALEV™
foscarnidopa/foslevodopa
Injection for subcutaneous use
12 mg/240 mg per mL



Plan it out by cleaning your hands and workspace and gathering and checking your supplies

Prepare the syringe by adding the medication into it, then inserting it into the pump

Prime the tubing by filling it with medication

Place the cannula on clean skin and connect the tubing

Pump the medication by starting the pump



“It is effortless to me and just part of my daily routine.”

—Janice, a VYALEV patient*

*Janice is an actual VYALEV patient and was on prescribed therapy when she provided a testimonial. Changes in therapy status may have occurred since that time.

SELECT SAFETY INFORMATION

Seeing, hearing, or feeling things that are not real (hallucination), infusion site reactions and infections, and new or worsened uncontrolled sudden movements (dyskinesia) are common and sometimes serious side effects.

Please see **Full Important Safety Information on pages 12-13** and **Full Prescribing Information, including Medication Guide.**

Please see **Full Instructions for Use.**



VYALEV™
foscarbidopa/foslevodopa
Injection for subcutaneous use
12 mg/240 mg per mL

Support you need,
when you need it

VYALEV™ COMPLETE

VYALEV[™] COMPLETE

VYALEV[™]
foscarnidopa/foslevodopa
Injection for subcutaneous use
12 mg/240 mg per mL

Caring and personalized support

If you and your doctor decide VYALEV is right for you, you will have one-on-one support every step of the way with VYALEV Complete. Your dedicated **Nurse Ambassador*** can:



Discuss your insurance coverage



Prepare you for starting
VYALEV at your doctor's office



Practice with you until you
are prepared to administer
your treatment at home



Answer your questions
along the way



VYALEV Complete 24/7 Hotline[†]


If you have questions about VYALEV at any time, give us a call at **1-866-489-2538**

*Nurse Ambassadors are provided by AbbVie and do not work under the direction of your healthcare professional or give medical advice. They are trained to direct patients to their HCP for treatment-related advice, including further referrals.

[†]The 24/7 Hotline is staffed by nurses who are specifically trained on VYALEV and can be reached 24 hours per day, 7 days a week. After business hours, you may be asked to leave a message for a return call within 30 minutes.

Please see Full Important Safety Information on pages 12-13 and Full Prescribing Information, including Medication Guide.

Important Safety Information and Use



Use

VYALEV is a prescription medicine used for treatment of advanced Parkinson's disease in adults. VYALEV contains two medicines, foscarbidopa and foslevodopa.

Important Safety Information

What is the most important safety information I should know about VYALEV™ (foscarbidopa/foslevodopa)?


Do not take VYALEV if you currently take or have recently taken (within the last 14 days) a medication for depression called a nonselective monoamine oxidase (MAO) inhibitor. Ask your healthcare provider or pharmacist if you are not sure if you take an MAO inhibitor.

Tell your healthcare provider about all your medical conditions and the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. When used together, VYALEV and certain other medicines, including medications for high blood pressure, MAO inhibitors, antipsychotics, metoclopramide, and isoniazid, may affect each other and cause serious side effects.

VYALEV may cause other serious side effects. Talk to your healthcare provider before starting VYALEV and while on VYALEV if you have had or have any of the following:

- **Falling asleep without warning during normal daily activities.** VYALEV may cause you to fall asleep while you are doing daily activities, such as driving, which may result in an accident. This can happen as late as 1 year after you start VYALEV. Do not drive or operate machinery until you know how VYALEV affects you. Tell your healthcare provider if you take medicines that can make you sleepy, such as sleep medicines, antidepressants, or antipsychotics.
- **Seeing, hearing, or feeling things that are not real (hallucinations). This is a common and sometimes serious side effect.**
- **Unusual urges.** Some people taking medicines for Parkinson's disease, including VYALEV, have reported problems, such as gambling, compulsive eating, compulsive shopping, and increased sex drive.

Important Safety Information and Use



- **Infusion Site Reactions and Infections.** Some people using VYALEV have had reactions and infections at the infusion site. **This is a common and sometimes serious side effect.** Remove your cannula and **call your healthcare provider if you have any of the following symptoms of an infection:** local spreading of redness, pain, swelling, warmth, change in color when pressing area, or fever. Take oral carbidopa/levodopa tablets until you are able to resume VYALEV.
- **Uncontrolled sudden movements (dyskinesia). This is a common and sometimes serious side effect.** If you have new dyskinesia or your dyskinesia gets worse, tell your healthcare provider. This may be a sign that your dose of VYALEV or other Parkinson's medicines may need to be adjusted.
- **Heart attack or other heart problems.** Tell your healthcare provider if you have had increased blood pressure, a fast or irregular heartbeat, or chest pain.
- Worsening of the **increased pressure in your eyes** (glaucoma). The pressure in your eyes should be checked after starting VYALEV.

Do not stop using VYALEV or change your dose unless you are told to do so by your healthcare provider. Tell your healthcare provider if you develop withdrawal symptoms, such as fever, confusion, or severe muscle stiffness.


These are not all the possible side effects of VYALEV. For more information, ask your healthcare provider or pharmacist.

VYALEV (foscarbidopa and foslevodopa) injection for subcutaneous use is available in a 120 mg foscarbidopa and 2,400 mg foslevodopa per 10 mL (12 mg foscarbidopa and 240 mg foslevodopa per mL) solution.

Please see the full Prescribing Information, including Medication Guide, for additional information about VYALEV. Talk to your healthcare provider if you have questions.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

If you are having difficulty paying for your medicine, AbbVie may be able to help. Visit AbbVie.com/PatientAccessSupport to learn more.



VYALEV™
foscarbidopa/foslevodopa
Injection for subcutaneous use
12 mg/240 mg per mL

Steps you can take **today**:



Ready to get started with VYALEV?

Talk to your doctor about enrolling.



Want to learn more about VYALEV?

Register for a Patient Education Program.

VYALEV **Patient Education** **Program**

Call **1-866-489-2538**, or visit vylev.com/learn to sign up.

Please see Full Important Safety Information on pages 12-13 and Full Prescribing Information, including Medication Guide.